


AP Calculus AB


Name



The rate at which ants arrive at a picnic is modeled by the function A, where A(t) is measured in ants per minute and t is measured in minutes. The graph of A for  $0 \le t \le 30$  is shown in the figure above. How many ants arrive at the picnic during the time interval  $0 \le t \le 30$ ?











## Wrap Up Accumulation



The rate at which people arrive at a theater box office is modeled by the function B, where B(t) is measured in people per minute and t is measured in minutes. The graph of B for  $0 \le t \le 20$  is shown in the figure above. Which of the following is closest to the number of people that arrive at the box office during the time interval  $0 \le t \le 20$ ?




3. A particle with velocity at any time t given by  $v(t) = e^t$  moves in a straight line. How far does the particle move from t = 0 to t = 2?




AP Calculus AB

## Wrap Up Accumulation



 $\chi(t) = e^{t} + C$  $\chi(z) - \chi(0)$ 

4. Snow is falling at a rate of  $r(t)=2e^{-0.1t}$  inches per hour, where *t* is the time in hours since the beginning of the snowfall. Which of the following expressions gives the amount of snow, in inches, that falls from time t = 0 to time t = 5 hours?



 $\int_{0}^{\infty} r(t) dt = \frac{2}{2} e^{0.1t} \int_{0}^{\infty}$ = -20 e +20