# Lesson 14 – General and Standard Form

## Goal:

- Can graph equations in general or standard form.
- Can build an equation for a linear function in general or standard form.
- Can use applications to model situations in general or standard form.

## New Terminology:

- General Form
- Standard form

## Discuss: Graph the line

3x - 5y + 9 = 0

# What strategy are you using?

					6					
					5-					
					4-					
					3-					
					2-					
					1_					
		1	1							
-6	-5 -	4 -:	3 -2	2 -'	1 0	1 2	2 3	3 4	+ 5	5 6
-6 -	.5 -	4 -:	3 -2	2 -'	1 0	1 2	2 3	3 4	- 5	56
-6	-5 -	4 -:	3 -2	2 -'	1 0 1- 2-	1 2	2 3	3 4	1 5	5 6
-6 ·	.5	4 -:	3 -2	2 -'	1 0 1- 2- 3-	1 2	2 3	3 4	- 5	5 6
-6	-5	4 -:	3 -:	2 -'	1 0 	1 2	2 3	3 4	4 5	5 6
-6	-5	4 -:	3 -2	2 - '	1 0 1- 2- 3- 4- 5-	1 2	2 3	3 4	+ 5	5 6

## Why might the given form be nicer than slope-intercept?

### **Chapter 7 Linear Functions**

We know how that using slope-intercept form is very useful and natural since it is in function form; however, it relies that the slope is well defined which we saw last class is not always the case.

Therefore, the **GENERAL FORM** is the general way of writing the equation of a line:

$$Ax + By + C = 0$$

## Where $A \ge 0$ and $B, C \in \mathbb{R}$

You already know how to graph a line in this form by changing it to slope-intercept form, but we can very easily graph it using the intercepts.

## Example: Graph the lines

$$\begin{aligned} x - 2y + 4 &= 0\\ y + 3 &= 0 \end{aligned}$$



**Practice**: Graph the following lines and label the *x*, *y*-intercepts.



**Discuss**: Determine the general form of the line that has an *x*-intercept of 3 and a *y*-intercept of 7.



**Discuss:** Determine the general form of the line that passes through the point (3,1) and has a slope of  $\frac{5}{4}$ .



**Discuss**: Determine the general form of the line that passes through the points (1, -9) and (-10, 7).



Notice that when we are solving for the intercepts, we consistently move the constant to the right side of the equation. This is called **STANDARD FORM**:

$$Ax + By = C$$

Where  $A \ge 0$ 

This is basically the same thing as general form, it just has the constant move on the opposite side as the variables. Since general form allows us to use the intercepts effectively it is useful in applications that involve the combination of two things adding to a fixed total.

**Example**: For a Christmas work party we have \$210 to spend on gift cards that come in \$10 and \$25 varieties. Determine an equation for the number of each gift card that can be bought.

**Practice**: You decide to go on a no carb diet and get all your calories from protein and fat. Your diet recommends 1800 calories each day. If 1 g of protein is 4 calories and 1 g of fat is 9 calories, determine an equation for the amount of protein and fat you can eat each day.

Discuss: What are the domain and range of these equations?

Assigned Problems: 7.2 page 365 – 369 # 2-8, 10, 11, 13, 14, 16, 20	
the second se	
12, 15, 19	
Key Ideas on page 364	